

Backward iteration in the unit ball

Olena Ostapyuk

Department of Mathematics
Kansas State University

AMS Meeting, Albuquerque NM, Spring 2010

One-dimensional case

Forward iteration

Let f be analytic self-map of $\mathbb{D} = \{z : |z| < 1\}$

n -th iterate of f $f_n = \underbrace{f \circ \dots \circ f}_{n \text{ times}}$

By **Schwarz's lemma**, f is a contraction in the pseudo-hyperbolic metric

$$d(z, w) = \left| \frac{z - w}{1 - \bar{w}z} \right|$$

Theorem (Denjoy-Wolff)

If a self-map of the disk f is not an elliptic automorphism, then there exist a unique point $p \in \overline{\mathbb{D}}$ such that the sequence $f_n(z)$ converges uniformly on compact subsets to p .

if $p \in \mathbb{D}$, then $f(p) = p$ and $|f'(p)| < 1$

if $p \in \partial\mathbb{D}$, then $f(p) = p$ and $0 < f'(p) \leq 1$ in the sense of non-tangential limits

One-dimensional case

Forward iteration

Let f be analytic self-map of $\mathbb{D} = \{z : |z| < 1\}$

n -th iterate of f $f_n = \underbrace{f \circ \dots \circ f}_{n \text{ times}}$

By **Schwarz's lemma**, f is a contraction in the pseudo-hyperbolic metric

$$d(z, w) = \left| \frac{z - w}{1 - \bar{w}z} \right|$$

Theorem (Denjoy-Wolff)

If a self-map of the disk f is not an elliptic automorphism, then there exist a unique point $p \in \overline{\mathbb{D}}$ such that the sequence $f_n(z)$ converges uniformly on compact subsets to p .

if $p \in \mathbb{D}$, then $f(p) = p$ and $|f'(p)| < 1$

if $p \in \partial\mathbb{D}$, then $f(p) = p$ and $0 < f'(p) \leq 1$ in the sense of non-tangential limits

One-dimensional case

Forward iteration

Let f be analytic self-map of $\mathbb{D} = \{z : |z| < 1\}$

n -th iterate of f $f_n = \underbrace{f \circ \dots \circ f}_{n \text{ times}}$

By **Schwarz's lemma**, f is a contraction in the pseudo-hyperbolic metric

$$d(z, w) = \left| \frac{z - w}{1 - \bar{w}z} \right|$$

Theorem (Denjoy-Wolff)

If a self-map of the disk f is not an elliptic automorphism, then there exist a unique point $p \in \overline{\mathbb{D}}$ such that the sequence $f_n(z)$ converges uniformly on compact subsets to p .

if $p \in \mathbb{D}$, then $f(p) = p$ and $|f'(p)| < 1$

if $p \in \partial\mathbb{D}$, then $f(p) = p$ and $0 < f'(p) \leq 1$ in the sense of non-tangential limits

One-dimensional case

Forward iteration

Let f be analytic self-map of $\mathbb{D} = \{z : |z| < 1\}$

n -th iterate of f $f_n = \underbrace{f \circ \dots \circ f}_{n \text{ times}}$

By **Schwarz's lemma**, f is a contraction in the pseudo-hyperbolic metric

$$d(z, w) = \left| \frac{z - w}{1 - \overline{w}z} \right|$$

Theorem (Denjoy-Wolff)

If a self-map of the disk f is not an elliptic automorphism, then there exist a unique point $p \in \overline{\mathbb{D}}$ such that the sequence $f_n(z)$ converges uniformly on compact subsets to p .

if $p \in \mathbb{D}$, then $f(p) = p$ and $|f'(p)| < 1$

if $p \in \partial\mathbb{D}$, then $f(p) = p$ and $0 < f'(p) \leq 1$ in the sense of non-tangential limits

One-dimensional case

Forward iteration

Let f be analytic self-map of $\mathbb{D} = \{z : |z| < 1\}$

n -th iterate of f $f_n = \underbrace{f \circ \dots \circ f}_{n \text{ times}}$

By **Schwarz's lemma**, f is a contraction in the pseudo-hyperbolic metric

$$d(z, w) = \left| \frac{z - w}{1 - \overline{w}z} \right|$$

Theorem (Denjoy-Wolff)

If a self-map of the disk f is not an elliptic automorphism, then there exist a unique point $p \in \overline{\mathbb{D}}$ such that the sequence $f_n(z)$ converges uniformly on compact subsets to p .

if $p \in \mathbb{D}$, then $f(p) = p$ and $|f'(p)| < 1$

if $p \in \partial\mathbb{D}$, then $f(p) = p$ and $0 < f'(p) \leq 1$ in the sense of non-tangential limits

The point p is called the **Denjoy-Wolff point** of f .

Cases:

1. $p \in \mathbb{D}$ f is called elliptic

2. $p \in \partial\mathbb{D}$, $f'(p) < 1$ hyperbolic

3. $p \in \partial\mathbb{D}$, $f'(p) = 1$ parabolic


The point p is called the **Denjoy-Wolff point** of f .

Cases:

1. $p \in \mathbb{D}$ f is called elliptic

2. $p \in \partial\mathbb{D}$, $f'(p) < 1$ hyperbolic

3. $p \in \partial\mathbb{D}$, $f'(p) = 1$ parabolic



Elliptic


The point p is called the **Denjoy-Wolff point** of f .

Cases:

1. $p \in \mathbb{D}$ f is called elliptic

2. $p \in \partial\mathbb{D}$, $f'(p) < 1$ hyperbolic

3. $p \in \partial\mathbb{D}$, $f'(p) = 1$ parabolic



Hyperbolic


The point p is called the **Denjoy-Wolff point** of f .

Cases:

1. $p \in \mathbb{D}$ f is called elliptic

2. $p \in \partial\mathbb{D}$, $f'(p) < 1$ hyperbolic

3. $p \in \partial\mathbb{D}$, $f'(p) = 1$ parabolic



Parabolic


The point p is called the **Denjoy-Wolff point** of f .

Cases:


1. $p \in \mathbb{D}$ f is called elliptic

2. $p \in \partial\mathbb{D}$, $f'(p) < 1$ hyperbolic


3. $p \in \partial\mathbb{D}$, $f'(p) = 1$ parabolic



Elliptic



Hyperbolic



Parabolic

If $p \in \partial\mathbb{D}$, **Julia's lemma** holds for the point p , and multiplier $c = f'(p) \leq 1$:

$$\forall R > 0 \quad f(H(p, R)) \subseteq H(p, cR),$$

where $H(p, R)$ is a horocycle at $p \in \partial\mathbb{D}$ of radius R :

$$H(p, R) := \left\{ z \in \mathbb{D} : \frac{|p - z|^2}{1 - |z|^2} < R \right\}$$

If $p \in \partial\mathbb{D}$, **Julia's lemma** holds for the point p , and multiplier $c = f'(p) \leq 1$:

$$\forall R > 0 \quad f(H(p, R)) \subseteq H(p, cR),$$

where $H(p, R)$ is a horocycle at $p \in \partial\mathbb{D}$ of radius R :


$$H(p, R) := \left\{ z \in \mathbb{D} : \frac{|p - z|^2}{1 - |z|^2} < R \right\}$$

If $p \in \partial\mathbb{D}$, **Julia's lemma** holds for the point p , and multiplier $c = f'(p) \leq 1$:

$$\forall R > 0 \quad f(H(p, R)) \subseteq H(p, cR),$$

where $H(p, R)$ is a horocycle at $p \in \partial\mathbb{D}$ of radius R :

$$H(p, R) := \left\{ z \in \mathbb{D} : \frac{|p - z|^2}{1 - |z|^2} < R \right\}$$



Backward iteration

Backward-iteration sequence:

$$\{z_n\}_{n=0}^{\infty}, f(z_{n+1}) = z_n \text{ for } n = 0, 1, 2 \dots$$

The sequence $d(z_n, z_{n+1})$ is increasing, so we need a bound on the pseudo-hyperbolic step: $d(z_n, z_{n+1}) \leq a < 1$

Theorem (Poggi-Corradini, 2003)

Let $\{z_n\}_{n=0}^{\infty}$ be a backward-iteration sequence for analytic self-map of the disk f with bounded pseudo-hyperbolic step $d(z_n, z_{n+1}) \leq a < 1$.

Then:

1. $z_n \rightarrow q \in \partial\mathbb{D}$, and q is a fixed point with a well-defined multiplier $f'(q) < \infty$
2. If $q \neq p$, then q is a **boundary repelling fixed point (BRFP)** (i.e. $f'(q) > 1$). If $q = p$, f is of parabolic type.
3. When q is BRFP, the convergence $z_n \rightarrow q$ is non-tangential.
4. If $q = p$, then $w_n \rightarrow q$ tangentially.

Backward iteration

Backward-iteration sequence:

$$\{z_n\}_{n=0}^{\infty}, f(z_{n+1}) = z_n \text{ for } n = 0, 1, 2, \dots$$

The sequence $d(z_n, z_{n+1})$ is increasing, so we need a bound on the pseudo-hyperbolic step: $d(z_n, z_{n+1}) \leq a < 1$

Theorem (Poggi-Corradini, 2003)

Let $\{z_n\}_{n=0}^{\infty}$ be a backward-iteration sequence for analytic self-map of the disk f with bounded pseudo-hyperbolic step $d(z_n, z_{n+1}) \leq a < 1$.

Then:

1. $z_n \rightarrow q \in \partial\mathbb{D}$, and q is a fixed point with a well-defined multiplier $f'(q) < \infty$
2. If $q \neq p$, then q is a **boundary repelling fixed point (BRFP)** (i.e. $f'(q) > 1$). If $q = p$, f is of parabolic type.
3. When q is BRFP, the convergence $z_n \rightarrow q$ is non-tangential.
4. If $q = p$, then $w_n \rightarrow q$ tangentially.

Backward iteration

Backward-iteration sequence:

$$\{z_n\}_{n=0}^{\infty}, f(z_{n+1}) = z_n \text{ for } n = 0, 1, 2, \dots$$

The sequence $d(z_n, z_{n+1})$ is increasing, so we need a bound on the pseudo-hyperbolic step: $d(z_n, z_{n+1}) \leq a < 1$

Theorem (Poggi-Corradini, 2003)

Let $\{z_n\}_{n=0}^{\infty}$ be a backward-iteration sequence for analytic self-map of the disk f with bounded pseudo-hyperbolic step $d(z_n, z_{n+1}) \leq a < 1$.

Then:

1. $z_n \rightarrow q \in \partial\mathbb{D}$, and q is a fixed point with a well-defined multiplier $f'(q) < \infty$
2. If $q \neq p$, then q is a **boundary repelling fixed point (BRFP)** (i.e. $f'(q) > 1$). If $q = p$, f is of parabolic type.
3. When q is BRFP, the convergence $z_n \rightarrow q$ is non-tangential.
4. If $q = p$, then $w_n \rightarrow q$ tangentially.

Backward iteration

Backward-iteration sequence:

$$\{z_n\}_{n=0}^{\infty}, f(z_{n+1}) = z_n \text{ for } n = 0, 1, 2, \dots$$

The sequence $d(z_n, z_{n+1})$ is increasing, so we need a bound on the pseudo-hyperbolic step: $d(z_n, z_{n+1}) \leq a < 1$

Theorem (Poggi-Corradini, 2003)

Let $\{z_n\}_{n=0}^{\infty}$ be a backward-iteration sequence for analytic self-map of the disk f with bounded pseudo-hyperbolic step $d(z_n, z_{n+1}) \leq a < 1$.

Then:

1. $z_n \rightarrow q \in \partial\mathbb{D}$, and q is a fixed point with a well-defined multiplier $f'(q) < \infty$
2. If $q \neq p$, then q is a **boundary repelling fixed point (BRFP)** (i.e. $f'(q) > 1$). If $q = p$, f is of parabolic type.
3. When q is BRFP, the convergence $z_n \rightarrow q$ is non-tangential.
4. If $q = p$, then $w_n \rightarrow q$ tangentially.

Backward iteration

Backward-iteration sequence:

$$\{z_n\}_{n=0}^{\infty}, f(z_{n+1}) = z_n \text{ for } n = 0, 1, 2, \dots$$

The sequence $d(z_n, z_{n+1})$ is increasing, so we need a bound on the pseudo-hyperbolic step: $d(z_n, z_{n+1}) \leq a < 1$

Theorem (Poggi-Corradini, 2003)

Let $\{z_n\}_{n=0}^{\infty}$ be a backward-iteration sequence for analytic self-map of the disk f with bounded pseudo-hyperbolic step $d(z_n, z_{n+1}) \leq a < 1$.

Then:

1. $z_n \rightarrow q \in \partial\mathbb{D}$, and q is a fixed point with a well-defined multiplier $f'(q) < \infty$
2. If $q \neq p$, then q is a **boundary repelling fixed point (BRFP)** (i.e. $f'(q) > 1$). If $q = p$, f is of parabolic type.
3. When q is BRFP, the convergence $z_n \rightarrow q$ is non-tangential.
4. If $q = p$, then $w_n \rightarrow q$ tangentially.

Backward iteration

Backward-iteration sequence:

$$\{z_n\}_{n=0}^{\infty}, f(z_{n+1}) = z_n \text{ for } n = 0, 1, 2, \dots$$

The sequence $d(z_n, z_{n+1})$ is increasing, so we need a bound on the pseudo-hyperbolic step: $d(z_n, z_{n+1}) \leq a < 1$

Theorem (Poggi-Corradini, 2003)

Let $\{z_n\}_{n=0}^{\infty}$ be a backward-iteration sequence for analytic self-map of the disk f with bounded pseudo-hyperbolic step $d(z_n, z_{n+1}) \leq a < 1$.

Then:

1. $z_n \rightarrow q \in \partial\mathbb{D}$, and q is a fixed point with a well-defined multiplier $f'(q) < \infty$
2. If $q \neq p$, then q is a **boundary repelling fixed point (BRFP)** (i.e. $f'(q) > 1$). If $q = p$, f is of parabolic type.
3. When q is BRFP, the convergence $z_n \rightarrow q$ is non-tangential.
4. If $q = p$, then $w_n \rightarrow q$ tangentially.

Backward iteration

Backward-iteration sequence:

$$\{z_n\}_{n=0}^{\infty}, f(z_{n+1}) = z_n \text{ for } n = 0, 1, 2, \dots$$

The sequence $d(z_n, z_{n+1})$ is increasing, so we need a bound on the pseudo-hyperbolic step: $d(z_n, z_{n+1}) \leq a < 1$

Theorem (Poggi-Corradini, 2003)

Let $\{z_n\}_{n=0}^{\infty}$ be a backward-iteration sequence for analytic self-map of the disk f with bounded pseudo-hyperbolic step $d(z_n, z_{n+1}) \leq a < 1$.

Then:

1. $z_n \rightarrow q \in \partial\mathbb{D}$, and q is a fixed point with a well-defined multiplier $f'(q) < \infty$
2. If $q \neq p$, then q is a **boundary repelling fixed point (BRFP)** (i.e. $f'(q) > 1$). If $q = p$, f is of parabolic type.
3. When q is BRFP, the convergence $z_n \rightarrow q$ is non-tangential.
4. If $q = p$, then $w_n \rightarrow q$ tangentially.

Multi-dimensional case

$$\mathbb{C}^N, \text{ inner product } (Z, W) = \sum_{j=1}^N Z_j \overline{W}_j, \quad \|Z\|^2 = (Z, Z)$$

$$\text{Unit ball } \mathbb{B}^N = \{Z \in \mathbb{C}^N : \|Z\| < 1\}$$

Julia's lemma in \mathbb{B}^N

Let f be a holomorphic self-map of \mathbb{B}^N and $X \in \partial\mathbb{B}^N$ such that

$$\liminf_{Z \rightarrow X} \frac{1 - \|f(Z)\|}{1 - \|Z\|} = \alpha < \infty$$

Then there exists a unique $Y \in \partial\mathbb{B}^N$ such that $\forall R > 0$
 $f(H(X, R)) \subset H(Y, \alpha R)$.

Horosphere of center $X \in \partial\mathbb{B}^N$ and radius $R > 0$:

$$H(X, R) = \left\{ Z \in \mathbb{B}^N : \frac{|1 - (Z, X)|^2}{1 - \|Z\|^2} < R \right\}$$

Multi-dimensional case

$$\mathbb{C}^N, \text{ inner product } (Z, W) = \sum_{j=1}^N Z_j \overline{W}_j, \quad \|Z\|^2 = (Z, Z)$$

$$\text{Unit ball } \mathbb{B}^N = \{Z \in \mathbb{C}^N : \|Z\| < 1\}$$

Julia's lemma in \mathbb{B}^N

Let f be a holomorphic self-map of \mathbb{B}^N and $X \in \partial\mathbb{B}^N$ such that

$$\liminf_{Z \rightarrow X} \frac{1 - \|f(Z)\|}{1 - \|Z\|} = \alpha < \infty$$

Then there exists a unique $Y \in \partial\mathbb{B}^N$ such that $\forall R > 0$
 $f(H(X, R)) \subset H(Y, \alpha R)$.

Horosphere of center $X \in \partial\mathbb{B}^N$ and radius $R > 0$:

$$H(X, R) = \left\{ Z \in \mathbb{B}^N : \frac{|1 - (Z, X)|^2}{1 - \|Z\|^2} < R \right\}$$

Multi-dimensional case

$$\mathbb{C}^N, \text{ inner product } (Z, W) = \sum_{j=1}^N Z_j \overline{W}_j, \quad \|Z\|^2 = (Z, Z)$$

$$\text{Unit ball } \mathbb{B}^N = \{Z \in \mathbb{C}^N : \|Z\| < 1\}$$

Julia's lemma in \mathbb{B}^N

Let f be a holomorphic self-map of \mathbb{B}^N and $X \in \partial\mathbb{B}^N$ such that

$$\liminf_{Z \rightarrow X} \frac{1 - \|f(Z)\|}{1 - \|Z\|} = \alpha < \infty$$

Then there exists a unique $Y \in \partial\mathbb{B}^N$ such that $\forall R > 0$

$$f(H(X, R)) \subset H(Y, \alpha R).$$

Horosphere of center $X \in \partial\mathbb{B}^N$ and radius $R > 0$:

$$H(X, R) = \left\{ Z \in \mathbb{B}^N : \frac{|1 - (Z, X)|^2}{1 - \|Z\|^2} < R \right\}$$

Multi-dimensional case

$$\mathbb{C}^N, \text{ inner product } (Z, W) = \sum_{j=1}^N Z_j \overline{W}_j, \quad \|Z\|^2 = (Z, Z)$$

$$\text{Unit ball } \mathbb{B}^N = \{Z \in \mathbb{C}^N : \|Z\| < 1\}$$

Julia's lemma in \mathbb{B}^N

Let f be a holomorphic self-map of \mathbb{B}^N and $X \in \partial\mathbb{B}^N$ such that

$$\liminf_{Z \rightarrow X} \frac{1 - \|f(Z)\|}{1 - \|Z\|} = \alpha < \infty$$

Then there exists a unique $Y \in \partial\mathbb{B}^N$ such that $\forall R > 0$

$$f(H(X, R)) \subset H(Y, \alpha R).$$

Horosphere of center $X \in \partial\mathbb{B}^N$ and radius $R > 0$:

$$H(X, R) = \left\{ Z \in \mathbb{B}^N : \frac{|1 - (Z, X)|^2}{1 - \|Z\|^2} < R \right\}$$

Multi-dimensional version of Denjoy-Wolff theorem holds:

Theorem (MacCluer, 1983)

If f has no fixed points in \mathbb{B}^N , then f_n converges uniformly on compacta to $p \in \partial\mathbb{B}^N$, the number $c := \liminf_{Z \rightarrow p} \frac{1 - \|f(Z)\|}{1 - \|Z\|} \in (0, 1]$ is a multiplier of f at p .

f is called hyperbolic if $c < 1$ and parabolic if $c = 1$.

Siegel domain:

$$\mathbb{H}^N = \{(z, w) \in \mathbb{C} \times \mathbb{C}^{N-1} : \operatorname{Re} z > \|w\|^2\}$$

Cayley transform: $\mathcal{C} : \mathbb{B}^N \rightarrow \mathbb{H}^N$

$$\mathcal{C}((z, w)) = \left(\frac{1+z}{1-z}, \frac{w}{1-z} \right)$$

$$\mathcal{C}^{-1}((z, w)) = \left(\frac{z-1}{z+1}, \frac{2w}{z+1} \right)$$

Multi-dimensional version of Denjoy-Wolff theorem holds:

Theorem (MacCluer, 1983)

If f has no fixed points in \mathbb{B}^N , then f_n converges uniformly on compacta to $p \in \partial\mathbb{B}^N$, the number $c := \liminf_{Z \rightarrow p} \frac{1 - \|f(Z)\|}{1 - \|Z\|} \in (0, 1]$ is a multiplier of f at p .

f is called hyperbolic if $c < 1$ and parabolic if $c = 1$.

Siegel domain:

$$\mathbb{H}^N = \{(z, w) \in \mathbb{C} \times \mathbb{C}^{N-1} : \operatorname{Re} z > \|w\|^2\}$$

Cayley transform: $\mathcal{C} : \mathbb{B}^N \rightarrow \mathbb{H}^N$

$$\mathcal{C}((z, w)) = \left(\frac{1+z}{1-z}, \frac{w}{1-z} \right)$$

$$\mathcal{C}^{-1}((z, w)) = \left(\frac{z-1}{z+1}, \frac{2w}{z+1} \right)$$

Multi-dimensional version of Denjoy-Wolff theorem holds:

Theorem (MacCluer, 1983)

If f has no fixed points in \mathbb{B}^N , then f_n converges uniformly on compacta to $p \in \partial\mathbb{B}^N$, the number $c := \liminf_{Z \rightarrow p} \frac{1 - \|f(Z)\|}{1 - \|Z\|} \in (0, 1]$ is a multiplier of f at p .

f is called hyperbolic if $c < 1$ and parabolic if $c = 1$.

Siegel domain:

$$\mathbb{H}^N = \{(z, w) \in \mathbb{C} \times \mathbb{C}^{N-1} : \operatorname{Re} z > \|w\|^2\}$$

Cayley transform: $\mathcal{C} : \mathbb{B}^N \rightarrow \mathbb{H}^N$

$$\mathcal{C}((z, w)) = \left(\frac{1+z}{1-z}, \frac{w}{1-z} \right)$$

$$\mathcal{C}^{-1}((z, w)) = \left(\frac{z-1}{z+1}, \frac{2w}{z+1} \right)$$

Theorem 1.

Let f be a analytic self-map of \mathbb{B}^N of hyperbolic type (with Denjoy-Wolff point $p \in \partial\mathbb{B}^N$), $\{Z_n\}$ be a backward-iteration sequence with bounded pseudo-hyperbolic step $d_{\mathbb{B}^N}(Z_n, Z_{n+1}) \leq a < 1$. Then:

1. There exists a point $\partial\mathbb{B}^N \ni \tau \neq p$ such that $Z_n \xrightarrow[n \rightarrow \infty]{} \tau$
2. $\{Z_n\}$ stays in a Koranyi region with vertex τ
3. Julia's lemma holds for τ with multiplier $\alpha \geq \frac{1}{c}$, where c is the multiplier at p , i.e. $f(H(\tau, R)) \subset H(\tau, \alpha R) \forall R > 0$

Since $\alpha \geq \frac{1}{c} > 1$, the point $\tau \in \partial\mathbb{B}^N$ is called the **boundary repelling fixed point** (BRFP) for f .

Characterization of BRFP in terms of backward-iteration sequences: Every backward-iteration sequence with bounded hyperbolic step converges to a BRFP; and if BRFP is isolated, then we can construct a backward-iteration sequence with bounded hyperbolic step that converges to it.

Theorem 1.

Let f be a analytic self-map of \mathbb{B}^N of hyperbolic type (with Denjoy-Wolff point $p \in \partial\mathbb{B}^N$), $\{Z_n\}$ be a backward-iteration sequence with bounded pseudo-hyperbolic step $d_{\mathbb{B}^N}(Z_n, Z_{n+1}) \leq a < 1$. Then:

1. There exists a point $\partial\mathbb{B}^N \ni \tau \neq p$ such that $Z_n \xrightarrow[n \rightarrow \infty]{} \tau$
2. $\{Z_n\}$ stays in a Koranyi region with vertex τ
3. Julia's lemma holds for τ with multiplier $\alpha \geq \frac{1}{c}$, where c is the multiplier at p , i.e. $f(H(\tau, R)) \subset H(\tau, \alpha R) \forall R > 0$

Since $\alpha \geq \frac{1}{c} > 1$, the point $\tau \in \partial\mathbb{B}^N$ is called the **boundary repelling fixed point** (BRFP) for f .

Characterization of BRFP in terms of backward-iteration sequences: Every backward-iteration sequence with bounded hyperbolic step converges to a BRFP; and if BRFP is isolated, then we can construct a backward-iteration sequence with bounded hyperbolic step that converges to it.

Theorem 1.

Let f be a analytic self-map of \mathbb{B}^N of hyperbolic type (with Denjoy-Wolff point $p \in \partial\mathbb{B}^N$), $\{Z_n\}$ be a backward-iteration sequence with bounded pseudo-hyperbolic step $d_{\mathbb{B}^N}(Z_n, Z_{n+1}) \leq a < 1$. Then:

1. There exists a point $\partial\mathbb{B}^N \ni \tau \neq p$ such that $Z_n \xrightarrow[n \rightarrow \infty]{} \tau$
2. $\{Z_n\}$ stays in a Koranyi region with vertex τ
3. Julia's lemma holds for τ with multiplier $\alpha \geq \frac{1}{c}$, where c is the multiplier at p , i.e. $f(H(\tau, R)) \subset H(\tau, \alpha R) \forall R > 0$

Since $\alpha \geq \frac{1}{c} > 1$, the point $\tau \in \partial\mathbb{B}^N$ is called the **boundary repelling fixed point** (BRFP) for f .

Characterization of BRFP in terms of backward-iteration sequences: Every backward-iteration sequence with bounded hyperbolic step converges to a BRFP; and if BRFP is isolated, then we can construct a backward-iteration sequence with bounded hyperbolic step that converges to it.

Theorem 1.

Let f be a analytic self-map of \mathbb{B}^N of hyperbolic type (with Denjoy-Wolff point $p \in \partial\mathbb{B}^N$), $\{Z_n\}$ be a backward-iteration sequence with bounded pseudo-hyperbolic step $d_{\mathbb{B}^N}(Z_n, Z_{n+1}) \leq a < 1$. Then:

1. There exists a point $\partial\mathbb{B}^N \ni \tau \neq p$ such that $Z_n \xrightarrow[n \rightarrow \infty]{} \tau$
2. $\{Z_n\}$ stays in a Koranyi region with vertex τ
3. Julia's lemma holds for τ with multiplier $\alpha \geq \frac{1}{c}$, where c is the multiplier at p , i.e. $f(H(\tau, R)) \subset H(\tau, \alpha R) \forall R > 0$

Since $\alpha \geq \frac{1}{c} > 1$, the point $\tau \in \partial\mathbb{B}^N$ is called the **boundary repelling fixed point** (BRFP) for f .

Characterization of BRFP in terms of backward-iteration sequences: Every backward-iteration sequence with bounded hyperbolic step converges to a BRFP; and if BRFP is isolated, then we can construct a backward-iteration sequence with bounded hyperbolic step that converges to it.

Theorem 1.

Let f be a analytic self-map of \mathbb{B}^N of hyperbolic type (with Denjoy-Wolff point $p \in \partial\mathbb{B}^N$), $\{Z_n\}$ be a backward-iteration sequence with bounded pseudo-hyperbolic step $d_{\mathbb{B}^N}(Z_n, Z_{n+1}) \leq a < 1$. Then:

1. There exists a point $\partial\mathbb{B}^N \ni \tau \neq p$ such that $Z_n \xrightarrow[n \rightarrow \infty]{} \tau$
2. $\{Z_n\}$ stays in a Koranyi region with vertex τ
3. Julia's lemma holds for τ with multiplier $\alpha \geq \frac{1}{c}$, where c is the multiplier at p , i.e. $f(H(\tau, R)) \subset H(\tau, \alpha R) \forall R > 0$

Since $\alpha \geq \frac{1}{c} > 1$, the point $\tau \in \partial\mathbb{B}^N$ is called the **boundary repelling fixed point** (BRFP) for f .

Characterization of BRFP in terms of backward-iteration sequences: Every backward-iteration sequence with bounded hyperbolic step converges to a BRFP; and if BRFP is isolated, then we can construct a backward-iteration sequence with bounded hyperbolic step that converges to it.

Theorem 1.

Let f be a analytic self-map of \mathbb{B}^N of hyperbolic type (with Denjoy-Wolff point $p \in \partial\mathbb{B}^N$), $\{Z_n\}$ be a backward-iteration sequence with bounded pseudo-hyperbolic step $d_{\mathbb{B}^N}(Z_n, Z_{n+1}) \leq a < 1$. Then:

1. There exists a point $\partial\mathbb{B}^N \ni \tau \neq p$ such that $Z_n \xrightarrow[n \rightarrow \infty]{} \tau$
2. $\{Z_n\}$ stays in a Koranyi region with vertex τ
3. Julia's lemma holds for τ with multiplier $\alpha \geq \frac{1}{c}$, where c is the multiplier at p , i.e. $f(H(\tau, R)) \subset H(\tau, \alpha R) \forall R > 0$

Since $\alpha \geq \frac{1}{c} > 1$, the point $\tau \in \partial\mathbb{B}^N$ is called the **boundary repelling fixed point** (BRFP) for f .

Characterization of BRFP in terms of backward-iteration sequences: Every backward-iteration sequence with bounded hyperbolic step converges to a BRFP; and if BRFP is isolated, then we can construct a backward-iteration sequence with bounded hyperbolic step that converges to it.

Problem:

Unlike in 1-dimensional case, not all BRFP's are isolated

Counterexample: $f : \mathbb{H}^2 \rightarrow \mathbb{H}^2, f(z, w) = (2z + w^2, w)$

Set of BRFP's: $\{(r^2, ir) \mid r \in \mathbb{R}\}$

Problem:

Unlike in 1-dimensional case, not all BRFP's are isolated

Counterexample: $f : \mathbb{H}^2 \rightarrow \mathbb{H}^2$, $f(z, w) = (2z + w^2, w)$


Set of BRFP's: $\{(r^2, ir) \mid r \in \mathbb{R}\}$

Problem:

Unlike in 1-dimensional case, not all BRFP's are isolated

Counterexample: $f : \mathbb{H}^2 \rightarrow \mathbb{H}^2$, $f(z, w) = (2z + w^2, w)$

Set of BRFP's: $\{(r^2, ir) \mid r \in \mathbb{R}\}$



Conjugations

1-dimensional hyperbolic case, forward iteration (Valiron, 1931):

$$\psi \circ f = \frac{1}{c}\psi,$$

where $\psi : \mathbb{D} \rightarrow \mathbb{H}$ is an analytic map to a half-plane.

1-dimensional case, backward iteration (Poggi-Corradini, 2000):

f an analytic self-map of \mathbb{D} with BRFP $1 \in \partial\mathbb{D}$ and multiplier α at 1 can be conjugated to the automorphism $\eta(z) = (z - a)/(1 - az)$, where $a = (\alpha - 1)/(\alpha + 1)$:

$$\psi \circ \eta(z) = f \circ \psi(z),$$

via an analytic map ψ of \mathbb{D} with $\psi(\mathbb{D}) \subseteq \mathbb{D}$, which has non-tangential limit 1 at 1.

Conjugations

1-dimensional hyperbolic case, forward iteration (Valiron, 1931):

$$\psi \circ f = \frac{1}{c}\psi,$$

where $\psi : \mathbb{D} \rightarrow \mathbb{H}$ is an analytic map to a half-plane.

1-dimensional case, backward iteration (Poggi-Corradini, 2000):

f an analytic self-map of \mathbb{D} with BRFP $1 \in \partial\mathbb{D}$ and multiplier α at 1 can be conjugated to the automorphism $\eta(z) = (z - a)/(1 - az)$, where $a = (\alpha - 1)/(\alpha + 1)$:

$$\psi \circ \eta(z) = f \circ \psi(z),$$

via an analytic map ψ of \mathbb{D} with $\psi(\mathbb{D}) \subseteq \mathbb{D}$, which has non-tangential limit 1 at 1.

Conjugations

N-dimensional case, forward iteration (Bracci, Gentili, Poggi-Corradini):

conjugation to a multiplication via $\psi : \mathbb{B}^N \rightarrow \mathbb{H}$:

$$\psi \circ f = \frac{1}{c}\psi$$

N-dimensional case, forward iteration (Bracci, Gentili, 2005):

f is conjugated to its linear part A via $\psi : \mathbb{H}^N \rightarrow \mathbb{H}^N$, assuming some regularity at the Denjoy-Wolff point:

$$\psi \circ f = A\psi$$

Conjugations

N-dimensional case, forward iteration (Bracci, Gentili, Poggi-Corradini):

conjugation to a multiplication via $\psi : \mathbb{B}^N \rightarrow \mathbb{H}$:

$$\psi \circ f = \frac{1}{c}\psi$$

N-dimensional case, forward iteration (Bracci, Gentili, 2005):

f is conjugated to its linear part A via $\psi : \mathbb{H}^N \rightarrow \mathbb{H}^N$, assuming some regularity at the Denjoy-Wolff point:

$$\psi \circ f = A\psi$$

Theorem 2. (N-dimensional case, backward iteration)

Suppose $f : \mathbb{H}^N \rightarrow \mathbb{H}^N$ is an analytic function and 0 is an isolated boundary repelling fixed point for f with multiplier $1 < \alpha < \infty$. Then f is conjugated to the automorphism $\eta(z, w) = (\alpha z, \sqrt{\alpha} w)$

$$\psi \circ \eta(Z) = f \circ \psi(Z),$$

via an analytic intertwining map ψ .

Construction of ψ :

$$\psi = \lim_{n \rightarrow \infty} \{f_n \circ \tau_n \circ p_1\}$$

where $p_1(z, w) := (z, 0)$ is the projection on the first (radial) dimension, so

$$\psi(z, w) = \psi(z, 0)$$

and is essentially one-dimensional map.

Theorem 2. (N-dimensional case, backward iteration)

Suppose $f : \mathbb{H}^N \rightarrow \mathbb{H}^N$ is an analytic function and 0 is an isolated boundary repelling fixed point for f with multiplier $1 < \alpha < \infty$. Then f is conjugated to the automorphism $\eta(z, w) = (\alpha z, \sqrt{\alpha} w)$

$$\psi \circ \eta(Z) = f \circ \psi(Z),$$

via an analytic intertwining map ψ .

Construction of ψ :

$$\psi = \lim_{n \rightarrow \infty} \{f_n \circ \tau_n \circ p_1\}$$

where $p_1(z, w) := (z, 0)$ is the projection on the first (radial) dimension, so

$$\psi(z, w) = \psi(z, 0)$$

and is essentially one-dimensional map.

Theorem 2. (N-dimensional case, backward iteration)

Suppose $f : \mathbb{H}^N \rightarrow \mathbb{H}^N$ is an analytic function and 0 is an isolated boundary repelling fixed point for f with multiplier $1 < \alpha < \infty$. Then f is conjugated to the automorphism $\eta(z, w) = (\alpha z, \sqrt{\alpha} w)$

$$\psi \circ \eta(Z) = f \circ \psi(Z),$$

via an analytic intertwining map ψ .

Construction of ψ :


$$\psi = \lim_{n \rightarrow \infty} \{f_n \circ \tau_n \circ p_1\}$$

where $p_1(z, w) := (z, 0)$ is the projection on the first (radial) dimension, so

$$\psi(z, w) = \psi(z, 0)$$

and is essentially one-dimensional map.

The image of ψ in \mathbb{H}^N :



Conjugation for expandable maps

Theorem 3.

Under some regularity condition, it is possible to improve ψ such that

$$\psi(z, w) = \psi(p_L(z, w)),$$

where p_L is a projection on the first L dimensions.

Condition is

$$f(z, w) = (\alpha z + o(|z|), Aw + o(|z|^{1/2}))$$

e.g. $A = \text{Diag}(\sqrt{\alpha}, \dots, \sqrt{\alpha}, \beta_1, \dots, \beta_{N-L})$, where $\beta_j < \sqrt{\alpha}$

Conjugation for expandable maps

Theorem 3.

Under some regularity condition, it is possible to improve ψ such that

$$\psi(z, w) = \psi(p_L(z, w)),$$

where p_L is a projection on the first L dimensions.

Condition is

$$f(z, w) = (\alpha z + o(|z|), Aw + o(|z|^{1/2}))$$

e.g. $A = \text{Diag}(\sqrt{\alpha}, \dots, \sqrt{\alpha}, \beta_1, \dots, \beta_{N-L})$, where $\beta_j < \sqrt{\alpha}$

Future goals

- Dimension of stable set at the BRFP q (union of all backward iteration sequences with bounded step tending to q)
- Conjugation for non-isolated fixed points
- Elliptic and parabolic cases

Thank you!

Future goals

- Dimension of stable set at the BRFP q (union of all backward iteration sequences with bounded step tending to q)
- Conjugation for non-isolated fixed points
- Elliptic and parabolic cases

Thank you!

Future goals

- Dimension of stable set at the BRFP q (union of all backward iteration sequences with bounded step tending to q)
- Conjugation for non-isolated fixed points
- Elliptic and parabolic cases

Thank you!

Future goals

- Dimension of stable set at the BRFP q (union of all backward iteration sequences with bounded step tending to q)
- Conjugation for non-isolated fixed points
- Elliptic and parabolic cases

Thank you!